

09/905,704

AMENDMENTS

provisional  
60/071,778  
02-27-07

In the Claims:

Please add claims 1-30 as follows:

1. An *in vivo* assay system for determining the effect of a pharmaceutically acceptable compound on angiogenesis comprising:

- a) a composition of microvascular endothelial cells; and
- b) a non-human host,

wherein said cells have a recombinant expression cassette encoding telomerase, and wherein said compound modulates the formation of functional microvessels from said cells that communicate with the circulatory system of said host.

OK abe  
change

2. The *in vivo* assay system of claim 1 further comprising a digital imaging device.

3. The *in vivo* assay system of claim 2 wherein said device detects fluorescence.

Thermon

4. The *in vivo* assay system of claim 1 wherein said cells stably express a transformed genetic marker.

Thermon

5. The *in vivo* assay system of claim 4 wherein said transformed genetic marker is enhanced green fluorescent protein (eGFP).

yang

6. The *in vivo* assay system of claim 1 wherein said cells are human dermal microvascular endothelial cells.

Thermon

7. The *in vivo* assay system of claim 1 wherein said telomerase is a human telomerase reverse transcriptase catalytic subunit.

scope to  
said  
mouse?

8. The *in vivo* assay system of claim 1 wherein said host is a SCID mouse.

9. The *in vivo* assay system of claim 1 wherein said compound is selected from the group consisting of growth factors, extracellular matrix molecules, proteinase inhibitors, cell adhesion molecules, angiostatic factors, apoptotic inducers, and inflammatory mediators.

10. The *in vivo* assay system of claim 9 wherein said compound is a growth factor.

11. The *in vivo* assay system of claim 10 wherein said growth factor is selected from the group consisting of angiopoietins, CTGF, EGF, FGF-2, IGF, PLGF, PDGF, SF, TGF, and VEGF.

12. The *in vivo* assay system of claim 11 wherein said growth factor is VEGF.

13. The *in vivo* assay system of claim 11 wherein said growth factor is FGF-2.

14. The *in vivo* assay system of claim 1 wherein said compound modulates tumor angiogenesis.

15. An *in vivo* method for analyzing the effect of a pharmaceutically acceptable compound on angiogenesis comprising:

a) providing a composition of microvascular endothelial cells, wherein said cells have a recombinant expression cassette encoding telomerase and a stably transformed genetic marker;

b) adding a compound that modulates the formation of functional microvessels to said cells to form a graft;

c) implanting said graft in a non-human host; and

d) determining the amount of neovascularization in said graft by measuring the expression of said transformed genetic marker.

16. The *in vivo* method of claim 15 wherein said cells are human dermal microvascular endothelial cells.

Thomas  
17. The *in vivo* method of claim 15 wherein said telomerase is a human telomerase reverse transcriptase catalytic subunit.

OKabe  
18. The *in vivo* method of claim 15 wherein said transformed genetic marker is enhanced green fluorescent protein (eGFP).  
don't use Thomas bc. although put GFP in cells did not use as a marker

OKabe  
19. The *in vivo* method of claim 15 wherein expression of said transformed genetic marker is detected by a digital imaging device.

Thomas  
20. The *in vivo* method of claim 15 wherein said compound is selected from the group consisting of growth factors, extracellular matrix molecules, proteinase inhibitors, cell adhesion molecules, angiostatic factors, apoptotic inducers, and inflammatory mediators.

Thomas  
21. The *in vivo* method of claim 20 wherein said compound is a growth factor.

Prewett  
22. The *in vivo* method of claim 21 wherein said compound is VEGF.

23. The *in vivo* method of claim 21 wherein said compound is FGF-2.

Bent  
24. The *in vivo* method of claim 15 wherein said composition further comprises matrigel.

Thomas  
25. The *in vivo* method of claim 15 wherein said host is a SCID mouse.

Prewett  
26. The *in vivo* method of claim 15 wherein said compound modulates tumor angiogenesis.

27. An *in vivo* system for human microvasculature formation comprising:  
a) a non-human host; and

b) at least one microvessel formed from a composition of microvascular endothelial cells having a recombinant expression cassette encoding telomerase and a stably transformed genetic marker in said host, wherein host blood is transmitted through said at least one microvessel.

28. The *in vivo* system of claim 27 wherein said host is a SCID mouse.

29. The *in vivo* system of claim 27 wherein said telomerase is a human telomerase reverse transcriptase catalytic subunit.

30. The *in vivo* system of claim 27 wherein said stably transformed genetic marker is enhanced green fluorescent protein (eGFP).